Finger Fill Canal Bibliography

Adkins, G. and P. Bowman. 1976. A study of the fauna in dredged canals of coastal Louisiana. Louisiana Wildlife and Fisheries Commission, Tech. Bull. 18. 72 p.

Arnold, EL Jr (1964): Man's alteration of estuaries by dredging and filling a grave threat to marine resources. Proc. Ann. Conf. S.E. Assoc. Game Fish Comm. 18, 269-273.

Bales, J. D. Fulford J. M. Swain E. *Review of Selected Features of the Natural System Model, and Suggestions for Applications in South Florida*, Water Resources Investigations Report 97-4039.

Bearden, D; Chung, KW; Strozier, ED; DeLorenzo, M; Sivertsen, S; Dias, A; Sanders, M; Macauley, JM; et al. 2002. Toxicological Studies in Tropical Ecosystems: an ecotoxicological Risk Assessment of Pesticide Runoff in South Florida Estuarine Ecosystems. Journal of Agricultural and Food Chemistry [J. Agric. Food Chem.]. Vol. 50, no. 15, pp. 4400-4408. 17 Jul 2002.

Brewster-Wingard, GL. 2001. Shaping south Florida's unique coastal ecosystems: Natural and anthropogenic influences. Geological Society of America. Programs with Abstracts [Geol. Soc. Am. Programs Abstracts]. Vol. 33, no. 2, pp. A-64. Mar 2001.
Brockmann, F.W. 1974. Seasonality of fishes in a south Florida brackish canal. Fla. Sci. 37(2):65-70.

Browder, J. A., A. Dragovich, J. Tashiro, E. Coleman-Duffie, C. Foltz, and J. Zweifel. 1986. *A Comparison of Biological Abundances in Three Adjacent Bay Systems Downstream from the Golden Gate Estates Canal System*, NOAA Technical Memorandum NMFS-SEFC-185. U.S. Department of Commerce.

Browder, J. A., J. Tashiro, E. Coleman-Duffie, and A. Rosenthal. 1988. "Comparison of Ichthyoplankton Immigration Rates into Three Bay Systems of the Ten Thousand Islands Affected by the Golden Gate Canal System." NOAA/National Marine Fisheries Service, Miami, FL.

Browder, J.A., A. Dragovich, J. Tashiro, E. Coleman-Duffie, C. Foltz, and J. Zweifel. 1986. A comparison of biological abundances in three adjacent bay systems downstream from the Golden Gate Estates canal system. NOAA Tech. Mem., NMFS-SEFC-185. 26 p.

Carriger, JF; Rand, GM; Gardinali, PR; Perry, WB; Tompkins, MS; Fernandez, AM. 2006. Pesticides of Potential Ecological Concern in Sediment from South Florida Canals: An Ecological Risk Prioritization for Aquatic Arthropods. Soil and Sediment contamination [Soil Sed. Contam.]. Vol. 15, no. 1, pp. 21-45. 2006.

Chesher, R.H. Canal survey Florida Keys. Mar.Res.Found.Inc., POB 2328, Key West, FL 33040, USA
Christensen, B.A. and R.M. Snyder, R.M. 1978. Establishment of residental waterfront property by construction of canal systems in coastal wetlands. Problems and solutions.

Coastal zone '78. Symposium on technical, environmental, socioeconomic and regulatory aspects of coastal zone management. San Francisco, CA(USA). Mar 14-16, 1978.

Cohn, J.P. 1994. Restoring the Everglades. BioScience 44(9):579-583.

Connolly, R. M. 2003. Differences in trophodynamics of commercially important fish between artificial waterways and natural coastal wetlands. *Estuarine, Coastal and Shelf Science* 58, no. 4: 929-36.

Connolly, R.M. 2003. Differences in trophodynamics of commercially important fish between artificial waterways and natural coastal wetlands. Estuarine, Coastal and Shelf Science 58(4): 929-936.

Finkl, C.W. and R.H. Charlier. 2003. Sustainability of Subtropical Coastal Zones in Southeastern Florida: Challenges for Urbanized Coastal Environments Threatened by Development, Pollution, Water Supply, and Storm Hazards. Journal of Coastal Research 19(4): 934-943.

Fox, A.M., W.T. Haller, K.D. Getsinger. 1991. Factors that influence water exchange in spring-fed tidal canals. Estuaries. Vol. 14, no. 4, pp. 404-413. 1991.

Genereux, D. E. Slater. 1999. Water exchange between canals and surrounding aquifer and wetlands in the Southern Everglades, USA. Journal of Hydrology (Amsterdam) [J. Hydrol. (Amst.)]. Vol. 219, no. 3-4, pp. 153-168. 8 Jul 1999.

Griffin, D.W., C.h.J.I. Gibson, E.K. Lipp, K. Riley, J.H.I. Paul, and J.B. Rose. 1999. Detection of viral pathogens by reverse transcriptase PCR and of microbial indicators by standard methods in the canals of the Florida Keys. Applied and Environmental Microbiology 65(9): 4118-4125.

Hansen, W.G., G. Bitton, J.L. Fox and P.L. Brezonik. 1977. Hydrocarbon status in Florida real estate canals. Mar. Pollut. Bull. Vol. 8, no. 3, pp. 57-62. 1977.

Hendon, J.R., M.S. Peterson and B.H. Comyns. (2000). Spatio-temporal distribution of larval Gobiosoma bosc adjacent to natural and altered marsh-edge habitats of Mississippi coastal waters. Bulletin of Marine Science 66(1):143-156.

Herke, W. H. 1959. Comparison of the length-weight relationship of several species of fish from two different, but connected, habitats. IN *Southeastern Association of Game and Fish Commisioners Thirteenth Annual Conference*, 299-313.

Hicks, D.B., T.R. Cavinder. Finger-fill canal studies: Florida and North Carolina. Athens, Georgia: U.S. Environmental Protection Agency, 1975. 427 p.: ill. (Call QH 541.5 .M3 F5 – Mote)

Irlandi, E., S. Macia, and J. Serafy. 1997. Salinity reduction from freshwater canal discharge: effects on mortality and feeding of an urchin (*Lytechinus variegatus*) and a

gastropod (*Lithopoma tectum*). *Bulletin of Marine Science* 61, no. 3: 869-79.

Koch, M; Zhang, G. 1998. Numerical modeling and management of saltwater seepage from coastal brackish canals in southeast Florida. Environmental Coastal Regions. pp. 395-404. 1998.

Koebel, J.W.J. 1995. An historical perspective on the Kissimmee River restoration project. Restoration Ecology 3(3): 149-159.

Koebel, J.W.J., B.L. Jones, and D.A. Arrington. 1999. Restoration of the Kissimmee River, Florida: Water quality impacts from canal backfilling. Environmental Monitoring and Assessment 57(1): 85-107.

Koebel, JW Jr; Jones, BL; Arrington, DA. 1999. Restoration of the Kissimmee River, Florida: Water quality impacts from canal backfilling Environmental Monitoring and Assessment [Environ. Monit. Assess.]. Vol. 57, no. 1, pp. 85-107. 1 Jul 1999.

Kruczynski, W.L. 1999. Water quality in the Florida Keys: Sources, effects, and solutions. EPA 904-R-99-005.

Kushlan, J.A. 1986. Exotic fishes of the Everglades: a reconsideration of proven impact. Environ. Conserv. 13(1):67-69.

Langevin, C.D., M.T. Stewart, and C.M. Beaudoin. 1998. Effects of sea water canals on fresh water resources: An example from Big Pine Key, Florida. Ground Water 36(3): 503-513.

Lewis, R.R., III, P. Clark, W.K. Fehring, H.S. Greening, R. Johansson, and R.T. Paul. 1999. The rehabilitation of the Tampa Bay Estuary, Florida, USA as an example of successful integrated coastal management. pp. 129-130 In: Proceedings of an international workshop on the rehabilitation of degraded coastal systems, 19-24 January 1998. Special publication 20. Phuket Marine Biological Center. Phuket.

Lewis, R.R., III. 1976. Impact of dredging in the Tampa Bay estuary, 1976-1976. pp. 31-55 In: Time-stressed coastal environments: assessment and future action. Proceedings of 2nd Annual Conference of The Coastal Society, New Orleans, LA, November 17-20, 1976.

Lincoln Smith, M.P., P.M.H. Hawes, and F.J. Duque-Portugal. 1995. Spatial variability in the nekton of a canal estate in southern New South Wales, Australia, and its implications for estuarine management. Marine & Freshwater Research 46(4): 715-721.

Lindall, W. N., W. A. Fable Jr., and L. A. Collins. 1975. Additional studies of the fishes, macroinvertebrates, and hydrological conditions of upland canals in Tampa Bay, Florida. *Fishery Bulletin* 73, no. 1: 81-85.

Lindall, W.N., Jr., J.R. Hall, and C.H. Saloman. 1973. Fishes, macroinvertebrates, and hydrological conditions of upland canals in Tampa Bay, Florida. Fishery Bulletin 71(1):155-163.

Lindall,WH Jr; Fable,WA Jr; Collins,LA (1975): Additional studies of the fishes, macroinvertebrates, and hydrological conditions of upland canals in Tampa Bay, Florida. Fish. Bull., U.S. 73, 81-85.

Lindall,WN (1973): Alterations of estuaries of South Florida: a threat to its fish resources. Mar. Freshw. Res. 35(10), 26-33.

Lindall,WN Jr; Hall,JR; Saloman,CH (1972): Fishes, macroinvertebrates, and hydrological conditions of upland canals in Tampa Bay, Florida. Fish.Bull., U.S. 71, 155-163.

Lindall,WN; Saloman,CH (1977): Alteration and destruction of estuaries affecting fishery resources of the Gulf of Mexico. Mar. Freshw. Res. 39(9),1-7.

Lindall,WN; Trent,L (1975): Housing development canals in the coastal zone of the Gulf of Mexico: ecological consequences, regulations, and recommendations. Mar. Fish. Rev. 37(6), 19-24.

Luther, G. W. III, S. Ma, R. Trouwborst, B. Glazer, M. Blickley, R. W. Scarborough and M. G. Mensinger. 2004. The roles of anoxia, H2S and storm events in fish kills of dead end canals of Delaware Inland Bays. Estuaries 27(3): 551-560. (View Abstract)

Luther, G.W., S. Ma, R. Trouwborst, B. Glazer, M. Blickley, R.W. Scarborough, and M.G. Mensinger. 2004. The Roles of Anoxia, H sub(2)S, and Storm Events in Fish Kills of Dead-end Canals of Delaware Inland Bays. Estuaries 27(3): 551-560.

Maxsted, J.R. et al. 1997. The ecological condition of dead-end canals of the Delaware and Maryland Coastal Bays. Estuaries 20:319-327

Miles, C.J. and R.J. Pfeuffer. 1997. Pesticides in canals of south Florida. Archives of Environmental Contamination and Toxicology 32(4): 337-345.

Morrill, J.B., Cde. Narvaez, R. Foster, F.B. II Ayer, and E. Connor. A report on the hydrography and biology of two man-made canal systems -Heron Lagoon and Grand Canal - on Siesta Key, Sarasota County, Florida. New Coll., Div. Nat. Sci., Sarasota, FL 33578, USA

Morrison. 1989. Ecological Assessment of the Cape Coral Residential Waterway System – available from Cape Coral.

Morton, R. M. 1992. Fish assemblages in residential canal developments near the mouth of a subtropical Queensland estuary. *Australian Journal of Marine and Freshwater*

Research 43: 1359-71.

Morton, R.M. 1989. Hydrology and fish fauna of canal developments in an intensively modified Astralian estuary. Estuar. Coastal Shelf Sci. 28:43-58.

Morton, R.M. 1992. Fish assemblages in residential canal developments near the mouth of a subtropical Queensland estuary. Aust. J. Mar. Freshw. Res. 43:1359-1371.

Neill, C. and R.E. Turner. 1987. Comparison of fish communities in open and plugged backfilled canals in Louisiana coastal marshes. North American Journal of Fisheries Management 7:57-62.

Nico, L.G. 2005. Changes in the fish fauna of the Kissimmee River Basin, peninsular Florida: nonnative additions. pp. 523-556 In: J.N. Rinne, R.M. Hughes, and B. Calamusso (eds.). Historical changes in large river fish assemblages of the Americas. American Fisheries Society Symposium 45.

Paulson, O.L., Jr, G.F. Pessoney, L. Massey, and D. Weaver. Reconnaissance of the flushing characteristics and water quality in coastal canals of the Gulf of Mexico. Source? Univ.Southern Mississippi, Dep.Geol., Hattiesburg, MS 39402, USA

Peterson, M.S., B.H. Comyns, J.R. Hendon, P.A. Bond and G.A. Duff. (2000). Habitat use by early life-history stages of fishes and crustaceans along a changing estuarine landscape: differences between natural and altered shoreline sites. Wetlands Ecology and Management 8(2/3):209-219.

Roessler, M. A. 1970. Checklist of fishes in Buttonwood Canal, Everglades National Park, Florida, and observations on the seasonal occurrence and life histories of selected species. *Bulletin of Marine Science* 20: 860-889.

Rozas, L. P. l992. Comparison of nekton habitats associated with pipeline canals and natural channels in Lousiana salt marshes. *Wetlands* 12, no. 2: 136-46.

Rozas, L.P. and D.J. Reed. 1994. Comparing nekton assemblages of subtidal habitats in pipeline canals transversing brackish and saline marshes in coastal Louisiana. Wetlands 14(4):262-275.

Saenger, P., and C. C. McIvor. 1975. Water quality and fish populations in a mangrove estuary modified by residential canal developments. *The International Symposium on Biology and Management of Mangroves*G. E. Walsh, S. C. Snedaker, and H. J. Teas, 753-65. Proceedings of the International Symposium on Biology and Management of Mangroves, Vol. 2, Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida.

Scott, GI; Fulton, MH; Wirth, EF; Chandler, GT; Key, PB; Daugomah, JW;
Serafy, J. E., K. C. Lindeman, T. E. Hopkins, and J. S. Ault. 1997. Effects of freshwater canal discharge on fish assemblages in a subtropical bay: field and laboratory

observations. *Marine Ecology Progress Series* 160: 161-72.

Sikkema, D.A. and P.C. Rosendahl. 1981. Prediction and assessment of the hydrologic conditions following the restoration of flow within an altered riverine system. Wetlands 1: 105-111.

State of Florida Dept. of Pollution Control. 1973. Survey of WQ in Waterways and canals of the Florida Keys - available from Cape Coral.

Surge, DM; Lohmann, KC. 2002. Temporal and Spatial Differences in Salinity and Water Chemistry in SW Florida Estuaries: Effects of Human-impacted Watersheds. 2002. Estuaries [Estuaries]. Vol. 25, no. 3, pp. 393-408. Jun 2002.

SWFRPC. 1984. Alternatives to Vertical Bulkheads for Shoreline Stabilization - available from Cape Coral.

SWFRPC. 1984. Cape Coral 208 Water Quality Study - available from Cape Coral.
Waller, B. G. 1976. "Analysis of Selected Benthic Communities in the Florida Everglades with Reference to their Physical and Chemical Environment." Water Resources Investigations 76-28. U.S. Geological Survey, Miami, FL.

Warburton, K. 1978. Community structure, abundance and diversity of fish in a Mexican coastal lagoon system. *Estuarine and Coastal Marine Science* 7: 497-519.

Weinstein, M. P., C. M. Courtney, and J. C. Kinch. 1977. The Marco Island estuary: a summary of physicochemical and biological parameters. *Florida Scientist* 40, no. 2: 97-124.

Woodburn, K.D. 1965. A discussion and selected, annotated references of subjective or controversial marine matters. Fla. State Bd. Conserv., Mar. Lab., Tech. Ser. 46. 50 p.

Other Sources
Dave Burr
Cape Coral Neighborhoods Group